

# Laboratory exercise

## Mössbauer course

Taha Ahmed and Adam Sobkowiak


Department of Chemistry

March 16, 2012




UPPSALA  
UNIVERSITET


# Folded Mössbauer spectra




# First guess




## Second guess



# Third guess




## Third guess—likely correct peak assignment



# Peak assignment


Doublet	Velocity/ $\text{mm s}^{-1}$	CS & $ \text{QS} $	Likely iron-species
Blue peaks	-0.33	CS=1.25	$\text{Fe}^{2+}$ , hs (8CN)
	2.83	$ \text{QS} =3.16$	$\text{Fe}^{2+}$ , hs (4CNT, 6CN)
Green peaks	-0.09	CS=1.29	$\text{Fe}^{2+}$ , hs (8CN)
	2.67	$ \text{QS} =2.76$	$\text{Fe}^{2+}$ , hs (4CNT, 6CN)
Red peaks	0.16	CS=0.41	$\text{Fe}^{3+}$ , hs (6CN, 4CN)
	0.65	$ \text{QS} =0.49$	$\text{Fe}^{3+}$ , hs (6CN, 4CNT)

And then we were told it was . . .




## Mössbauer parameters from literature


Sample	Compound	CS/mm s <sup>-1</sup>	QS/mm s <sup>-1</sup>
natural	FeSO <sub>4</sub>	1.27 ± 0.02	2.71 ± 0.02
	FeSO <sub>4</sub>	1.25 ± 0.03	2.71 ± 0.03
	FeSO <sub>4</sub>	1.30 ± 0.05	2.65 ± 0.05
synthetic	Pure FeSO <sub>4</sub>	1.27 ± 0.01	2.72 ± 0.02




# Our assignment compared to literature





# Iron sulfate heptahydrate: $\text{Fe}^{\text{II}}(\text{SO}_4) \cdot 7 \text{H}_2\text{O}$



# Iron sulfate pentahydrate: $\text{Fe}^{\text{III}}(\text{SO}_4)_3 \cdot 5 \text{H}_2\text{O}$



# Iron sulfate nonahydrate: $\text{Fe}_2^{\text{III}}(\text{SO}_4)_3 \cdot 9\text{H}_2\text{O}$



## Bibliography

- [1] Tore Ericsson. *Mössbauer in Uppsala*. Uppsala: Uppsala universitet, Fysiska institutionen, 2000.
- [2] M. D. Lane et al. 'Mineralogy of the Paso Robles soils on Mars'. *American Mineralogist* 93.5-6 (May 2008), pp. 728–739. DOI: 10.2138/am.2008.2757.
- [3] Pedro A. Montano. 'Mössbauer spectroscopy of iron compounds found in West Virginia coals'. *Fuel* 56.4 (Oct. 1977), pp. 397–400. DOI: 10.1016/0016-2361(77)90066-7.